Research Article

ETHNOBOTANICAL STUDIES ON WILD EDIBLE PLANTS OF GOND, HALBA AND KAWAR TRIBES OF SALEKASA TALUKA, GONDIA DISTRICT, MAHARASHTRA STATE, INDIA

Patale Chandrakumar K ^{1*}, Nasare Praveenkumar N ², Narkhede Sushama D ¹

¹Department of Botany, Government Science College, Gadchiroli, Maharashtra, India

²Department of Botany, Nilkanthrao Shinde Science & Arts College, Bhadravati, Dist- Gadchiroli, Maharashtra, India

*Corresponding Author Email: ckpatle@gmail.com

Article Received on: 21/05/15 Revised on: 30/06/15 Approved for publication: 17/07/15

DOI: 10.7897/2230-8407.068103

ABSTRACT

The ethnobotanical studies on wild edible plant was conducted among the Gond, Halba and Kavar tribes of Salekasa taluka, Gondia district, Maharashtra state, India, through survey, interview and field work along with knowledgeable persons during December 2013 – May 2015. Total 80 wild edible plant species recorded which belongs to 69 genera and 38 Families. Among 38 families, 5 belong to monocot and rests 33 are of dicotyledons. The most widely utilized species belongs to Amaranthaceae (7), Araceae (7), Caesalpinaceae (6), Fabaceae (5) and Solanaceae (4). The families Euphorbiaceae, Poaceae, Rubiaceae, Rutaceae, Tiliaceae represented by 3 species each and remaining families represented by one or two species each. In the present study, total 80 species are documented; among them 31 are herbs, 5 shrubs, 10 climbers, 33 trees and 1 parasite.

Keywords: Ethnobotany, wild edible plants, Gond, Halba, Kavar Tribe, Salekasa taluka

INTRODUCTION

Ethnobotany is one of the most interesting themes of economic botany which might have first of all came into the existence probably when earliest man of “stone age” observed the animals mostly the apes and monkeys eating certain plants or plant parts ex. Fruits, leaves and even inflorescences to satisfy their hunger. Therefore, on the basis of plants usage first of all by animals and later on by the human beings the concepts of Ethnobotany and Ethnozoology were evolved, which merged into a common term known as Ethnobiology. However, the term Ethnobotany was first of all used in the last of 19th century by J. W. Harsh Berger (1895) to indicate the interrelationship of plants with aboriginal people or tribal societies ¹. In many parts of the world, wild plants are obtained from forests or wild areas designated for extractive resources and managed by local communities ². Wild edible plants provide food quantity as well as make significant contribution to the population nutrition throughout the year ^{3,4}. Wild fruits provide nutrition for the forest dwellers and many of marginalized rural communities ⁵. The contributions of forest foods security can be categorized into three main ways viz. i) providing a supplementary source of food, ii) as seasonal foods in the diet, and iii) as emergency food supplies during periods when others are unavailable ⁶. Survey and documentation of wild edible plants have been reported from several parts of India ⁷⁻³². The present studies were conducted to explore the knowledge of wild edible plants of Gond, Halba and Kavar tribe of Salekasa Taluka (Gondia district, Maharashtra state, India).

MATERIALS AND METHODS

Study area

Salekasa taluka belongs to Deori sub division of Gondia district in Nagpur revenue division in the Berar region in the state of Maharashtra, India. The Hazara Fall, Ambezaran, Daldali, Gendurzirra, Ranidoh, Kachargarh cave and Bewartola water reserve

are the natural attraction for tourist. Banjari, Bewartola, Bharritola, Chandsuraj, Chiglutola, Daldalkuhi, Dandari, Dhannegaon, Dumbartola, Jamakudo, Kopalgargh, Murkutdoh, Sikaritola, Tekatola, Thubrutola, Toyagondi and Vicharpur are the tribal villages. Villages are situated in dense forest area and on hills, known as “Darekasa Hill Range”, it is the part of Satpuda hill ranges (Figure 1).

People

Total Population of the Salekasa taluka is 90,679 as per the survey of census during 2011 by Indian Government. Out of which total scheduled tribes are 23,900. It means that 26.36% is the tribal population. The eastern part of taluka is covered by dense Forest. Most of the tribal people live in forest area in hamlets with their customs and rituals. The main occupation of tribal people of Salekasa taluka is farming and related works. Rice is main crop of this region. Gond, Halba and Kavar are three tribes live since long times. The God of these tribes is Budalpen or Budadev. The meaning of Pen is God. The God Budalpen is situated at Kachargarh cave.

Data collection

The study was conducted among the Gond, Halba and Kavar peoples of Salekasa taluka through survey; interviews and field works along with knowledgeable persons during December 2013 – May 2015. To collect traditional knowledge on wild edible plants, frequent discussions were made with local persons, including Mukhias, Patels, Tribal leaders, Farmers, Shepherds, housewives and children. Information was noted in field books. Field work was completed with tribal peoples. Plant specimens were collected with the permission of Maharashtra State Biodiversity Board. Collected specimens and photographs shown to peoples for local names and usage. These specimens were identified with help of floras [33-37]. The identified plants are arranged alphabetically with family names, local names and parts used (Table 1).

Table 1: Wild edible plants used by Tribals of study area

S.N	Plant Name	Family	Local Name	Part used
1	<i>Abrus precatorius</i> L.	Fabaceae	Gomchi, Gunja	Leaves
2	<i>Acacia nilotica</i> (L.) Willd.	Mimosaceae	Babul	Gum
3	<i>Aegle marmelos</i> Corr	Rutaceae	Bel	Fruit
4	<i>Alternanthera sessilis</i> DC	Amaranthaceae	Jibhkati	Leaves
5	<i>Amaranthus spinosus</i> L.	Amaranthaceae	Matbhaji	Leaves
6	<i>Amaranthus viridis</i> L.	Amaranthaceae	Khedabhaji	Leaves
7	<i>Amorphophalus bhandarensis</i> Yadav, Kahalkar and Bhuskute	Araceae	Gaivar	Leaves
8	<i>Amorphophalus paeonifolius</i> Dernst	Araceae	Jimikanda	Leaves
9	<i>Anogeissus latifolia</i> (Roxb. ex. DC) Wall ex Guill & Perr.	Combretaceae	Dhawnra	Gum
10	<i>Antidesma ghaesembilla</i> Gaertn	Euphorbiaceae	Jondhurli, Amti	Fruit
11	<i>Bambusa vulgaris</i> Schrad	Poaceae	Vadud, Bans	Shoot
12	<i>Bauhinia purpurea</i> Linn.	Caesalpiniaceae	Koilar	Leaves
13	<i>Bauhinia vahlii</i> Wight & Arn.	Caesalpiniaceae	Mahur	Seeds
14	<i>Bridelia retusa</i> Spreng	Euphorbiaceae	Kasai	Fruit
15	<i>Buchanania cochinchinesis</i> (Lour.) Almeida	Anacardiaceae	Sadeka, Char	Seeds
16	<i>Butea monosperma</i> (Lam.) Taub.	Fabaceae	Parsa, Palas	Flower
17	<i>Cajanus scarabaeoides</i> (L.) Du Petit-thou	Fabaceae	Ran- tur	Seeds
18	<i>Cardiospermum helicacabum</i> L.	Sapindaceae	Kaparpothi	Fruits
19	<i>Cassia fistula</i> L.	Caesalpiniaceae	Bahava	Flower
20	<i>Cassia tora</i> L.	Caesalpiniaceae	Chirota, Charota	Leaves
21	<i>Celosia argentea</i> L.	Amaranthaceae	Siliyari	Leaves
22	<i>Cissus quadrangularis</i> L.	Vitaceae	Hadijod	Shoot
23	<i>Colocasia esculenta</i> (L.) Schott	Araceae	Aaki, Kochaimati, ghuya	Leaves, petiole & tuber
24	<i>Commelina benghalensis</i> L.	Commelinaceae	Telka bhaji	Leaves
25	<i>Corchorus aestuans</i> L.	Tiliaceae	Chechbhaji	Leaves
26	<i>Cordia dichotoma</i> Forst f.	Boraginaceae	Selvat	Fruit
27	<i>Costus speciosus</i> (Koen) Sm.	Zingiberaceae	Kevkanda	Leaves, Tuber
28	<i>Cryptocoryne retorspiralis</i> (Roxb.) Kunth	Araceae	Pakanbhed	Leaves
29	<i>Curculigo orchoides</i> Gaertn	Hypoxidaceae	Kali musli	Tuber
30	<i>Dendrocalamus strictus</i> (Roxb.) Nees	Poaceae	Ranj, bamboo	shoots
31	<i>Dendrophloe falcata</i> (L.f.) Etting	Loranthaceae	Vanda , chipur	Fruit
32	<i>Dillenia pentagyna</i> Roxb.	Dilleniaceae	Ran- kel	Fruits
33	<i>Dioscorea bulbifera</i> L.	Dioscoreaceae	Dangkanda	Tuber, bulbils
34	<i>Dioscorea oppositifolia</i> L.	Dioscoreaceae	Bhaisdheti	Tuber, bulbils
35	<i>Diospyros melanoxylon</i> Roxb.	Ebenaceae	Tembhrun, Tendu	Fruit
36	<i>Diplocyclos palmatus</i> (L.) Jeffrey	Cucurbitaceae	Shivlingi	Leaves, fruit
37	<i>Ficus racemosa</i> L.	Moraceae	Umbar	Fruit
38	<i>Gardenia latifolia</i> Ait.	Rubiaceae	Ghogar	Fruit
39	<i>Gardenia resinifera</i> Roth	Rubiaceae	Dhikemalhi	Fruit, gum
40	<i>Gmelina arborea</i> Roxb.	Verbenaceae	Sivan	Fruit
41	<i>Grevia hirsuta</i> Vahl.	Tiliaceae	Gursukdi, kolhati	Fruit
42	<i>Grevia tilifolia</i> Vahl	Tiliaceae	Dhaman	Fruits
43	<i>Hemidesmus indicus</i> (L.) R. Br.	Periplocaceae	Khaprilbela	Tuber
44	<i>Holarrhena pubescens</i> (Buch-Ham.) Wall. ex. G. Don.	Apocynaceae	Kudva	Flower, Fruit
45	<i>Ipomoea aquatica</i> Lour	Convolvulaceae	Karmotabhaji	Leaves
46	<i>Kalanchoe pinnata</i> (Lam.) Pers.	Crassulaceae	Panphuti, Ghavpala	Leaves
47	<i>Lantana camara</i> L.	Verbenaceae	Bantulsi	Fruit
48	<i>Lasia spinosa</i> (L.) Thw	Araceae	Gongal -kanda	Leaves
49	<i>Limonia acidissima</i> L.	Rutaceae	Kawath	Fruit
50	<i>Madhuca longifolia</i> (Koen.) Mac. Var. <i>latifolia</i> (Rox.) Chevalier	Sapotaceae	Idukmada, Mahu, mahuva	Petals
51	<i>Mangifera indica</i> L.	Anacardiaceae	Marka, Aamba	Fruit
52	<i>Merremia gangetica</i> (L.) Cuf.	Convolvulaceae	Kukripota	Leaves
53	<i>Mimusops elengi</i> L.	Sapotaceae	Massor	Fruits
54	<i>Mucuna pruriens</i> (L.) DC	Fabaceae	Kavaskuri	Seeds
55	<i>Murraya koenigii</i> (L.) Spreng.	Rutaceae	Mithhalimb	Leaves
56	<i>Nelumbo nucifera</i> Gaertn.	Nelumbonaceae	Pavan	Rhizome, Seeds
57	<i>Nymphaea nouchali</i> Burm. f.	Nymphaeaceae	Kamal	Rhizome
58	<i>Nymphaea rubra</i> Roxb.	Nymphaeaceae	Lalkamal	Rhizome
59	<i>Oxalis psittacorum</i> (Willd.) Vahl	Oxalaceae	Bindranipor	Leaves
60	<i>Oryza rufipogon</i> Griff.	Poaceae	Karnga, Devdhan	Fruit
61	<i>Phoenix sylvestris</i> (L.) Roxb.	Araceae	Sindi, Chhindi	Fruit
62	<i>Phyllanthus emblica</i> L.	Euphorbiaceae	Nalli, Amla	Fruit
63	<i>Physalis minima</i> L.	Solanaceae	Chirpoti	Fruit
64	<i>Pueraria tuberosa</i> (Roxb ex Willd) DC	Fabaceae	Bhuikanda	Tuber
65	<i>Schleichera oleosa</i> (Lour.) O. Ken	Sapindaceae	Kusum	Seeds

66	<i>Semecarpus anacardium</i> L.f.	Anacardiaceae	Kohka, Bhelva	Thalamus
67	<i>Senna occidentalis</i> (L.) Link.	Caesapiniaceae	Rantarota	Seeds
68	<i>Smilax zeylanica</i> L.	Smilacaceae	Sherdera, Ramdatun	Fruits
69	<i>Solanum nigrum</i> L.	Solanaceae	Kamuni	Leaves, Fruit
70	<i>Solanum torvum</i> Sw.	Solanaceae	Ranbhatai	Fruit
71	<i>Solanum virginianum</i> L.	Solanaceae	Bhaskatiya	Fruit
72	<i>Strychnos potatorum</i> L.f.	Loganiaceae	Kariakval, Nirmali	Fruit
73	<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae	Jambhul	Fruit
74	<i>Tamarindus indica</i> L.	Caesalpinaceae	Sitta, Chich, Imli	Fruit
75	<i>Tamilnandia uliginosa</i> (Retz) Tirvengadam & sastre	Rubiaceae	Telpendhra	Fruit
76	<i>Terminalia bellirica</i> Roxb.	Combretaceae	Behda	Seeds
77	<i>Theriophonum minutum</i> (Willd) baill.	Araceae	Undirkani	Leaves
78	<i>Trapa natans</i> L. var. <i>bispinosa</i> (Roxb.) Makino	Trapaceae	Shingara	Fruit
79	<i>Ziziphus mauritiana</i> Lam.	Rhamnaceae	Renga, Boir, Ber	Fruit
80	<i>Ziziphus oenoplia</i> (L.) Mill.	Rhamnaceae	Mokaiya, Ironi, Yaroni	Fruit

Figure 1: Showing study area in Gondia district map

Figure 2: Wild edible plant species in different categories

Figure 3: Life forms of wild edible plants

RESULTS AND DISCUSSION

The data analysis shows that, Gond, Halba and Kavar tribes of study area possess a tremendous knowledge on the wild edible plants. Total 80 plant species from 69 genera and 38 families have been recorded as wild edible plants in study area (Table 1). Among 38 families 5 families belongs to monocotyledons and rest from the dicotyledons. Of which fruit ranked first with 33 species, leaves, seeds, tubers, flower, rhizome, shoot, gum and thalamus ranked next with 23, 8, 6, 3, 3, 3, 3 and 1 species respectively (Figure 2). The most widely utilized species belongs to Amaranthaceae (7), Araceae (7), Caesalpiniaceae (6), Fabaceae (5) and Solanaceae (4). The families Euphorbiaceae, Poaceae, Rubiaceae, Rutaceae, Tiliaceae represented by 3 species each and remaining families represented by one or two species each. In the present study, total 80 species are listed; among them 31 are herbs, 5 shrubs, 10 climbers, 33 trees and 1 parasite (Figure 3).

Most of the edible parts used as leaves, tubers and fruits are consumed after cooking (*Alternanthera sessilis*, *Amaranthus spinosus*, *Amaranthus viridis*, *Amorphophalus bhandarensis*, *Amorphophalus paeonifolius*, *Cassia tora*, *Celosia argentea*, *Colocasia esculenta*, *Commelina benghalensis*, *Corchorus aestuans*, *Costus speciosus*, *Cryptocoryne retrospiralis*, *Lasia spinosa*, *Merremia gangetica*, *Oryza rufipogon*, *Solanum virginianum*, *Theriophonum minutum*, *Bambusa vulgaris*, *Dendrocalamus strictus*, *Solanum torvum*, *Tamilnandia uliginosa*). Some of the edible parts are roasted (*Dioscorea bulbifera*, *Dioscorea oppositifolia*, *Nymphaea nouchali*, *Nymphaea rubra*, *Pueraria tuberosa*, *Trapa natans*). Some of the

plant parts are directly consumed as fresh (*Aegle marmelos*, *Antidesma ghaesembilla*, *Bridelia retusa*, *Dendrophthoe falcata*, *Grevia hirsuta*, *Hemidesmus indicus*, *Lantana camara*, *Madhuca longifolia*, *Semecarpus anacardium*, *Tamarindus indica*, *Terminalia bellirica*, *Ziziphus mauritiana*, *Zizyphus oenoplia*). Many plants products are stored after proper preparations and used all year around, some of them are *Amorphophalus paeonifolius*, *Colocasia esculenta*, *Cordia dichotoma*, *Dioscorea oppositifolia*, *Mangifera indica*, *Phyllanthus emblica*, *Pueraria tuberosa*, *Tamarindus indica*.

CONCLUSION

The Men and Women of Gond, Halba and Kavar tribes of study area are well experienced and have rich knowledge in utilizing of wild edible plants. This important knowledge is slowly diminishing day by day due to invasion of alien cultures. Today, it is urgent need to document the indigenous knowledge for future generations and also encourage these tribes for cultivation of wild edible plants in their home gardens.

ACKNOWLEDGEMENT

Authors are thanks to Dr. J. M. Khobragade, Principal, Government Science College, Gadchiroli and Dr. S. M. Bhuskute, Principal, Bhawbhuti Mahavidyalaya, Amgaon for valuable guidance and help. Authors are also thankful to the local, tribal people of the study area who have provided valuable information about the uses of plants and help in the collection of information.

Tamilnandia uliginosa

Cryptocoryne retrospiralis

Ipomoea aquatica

Celosia argentea

Phoenix sylvestris

Dendrophloe falcata

Butea monosperma

Amaranthus spinosus

Boiled tuber of *Dioscorea oppositifolia*

Costus speciosus

Cassia tora

Trapa natans var. *bispinosa*

Dillenia pentagyna

Smilax zeylanica

Buchanania cochinchinensis

Olax psittacorum

Nymphaea nouchali (Rhizome)

Amorphophalus paeonifolius

Diospyros melanoxylon

Cordia dichotoma

Ziziphus mauritiana

Solanum virginianum

Solanum torvum

Colocasia esculenta

Theriophonum minutum

Nymphaea nouchali

Children eating seeds of *Nymphaea nouchali*

REFERENCES

- Trivedi PC, Sharma N, Text book of Ethnobotany, Pointer publishers, Jaipur, India, 2011.
- Jadhav VD, Mahadkar SD, Valvi SR, Documentation and ethnobotanical survey of wild edible plants from Kolhapur district, Recent Research in Science and technology, 2011; 3 (12): 58-63.
- Grivetti LE, Ogle BM, Value of traditional foods in meeting macro and micronutrient need: the wild plant connection, Natl. Res. Rev, 2000; 3: 31-46. <http://dx.doi.org/10.1079/095442200108728990>
- Ogle BM, Ho TT, Hoang ND, Nguyen NXD, Food, Feed or Medicine: The multiple functions of edible wild plants in Vietnam, Economic botany, 2003; 57 (1): 103-117. [http://dx.doi.org/10.1663/0013-0001\(2003\)057\[0103:FFOMTM\]2.0.CO;2](http://dx.doi.org/10.1663/0013-0001(2003)057[0103:FFOMTM]2.0.CO;2)
- Nazurudeen A, Nutritional composition of some lesser known fruits used by the ethnic communities and local folks of Kerala, Indian Journal of traditional knowledge, 2010; 9 (2): 398-402.
- Rasingam L, Ethnobotanical studies on the wild edible plants of Irula tribes of Pillar Valley, Coimbatore district, Tamil Nadu, India, Asian Pacific Journal of Tropical Biomedicine, 2012, S1493- S1497. [http://dx.doi.org/10.1016/S2221-1691\(12\)60443-2](http://dx.doi.org/10.1016/S2221-1691(12)60443-2)
- Deb D, Sarkar A, Deb BB, Datta BK, Muzumdar K, Wild edible plants and their utilization in traditional recipes of Tripura, Northeast India, Advances in biological research, 2013; 7 (5): 203- 211.
- Kar A, Borthakur SK, Wild vegetables sold in markets of Karbi Anglong, Assam, Indian Journal of Traditional Knowledge, 2007; 6 (1): 169-172.
- Ballabh B, Chaurasia OP, Pande PC, Ahmed Z, Raw edible plants of cold desert ladakh, Indian Journal of Traditional Knowledge, 2007; 6 (1): 182-184.
- Misra S, Maikhuri RK, Kala CP, Rao KS, Saxena KG, Wild leafy vegetables: A study of their subsistence dietetic support to the inhabitants of Nanda Devi Biosphere Reserve, India, Journal of Ethnobiology and Ethnomedicine, 2008; 4 (15) : 1-9.
- Sasu R, Rajendran A, Maharajan M, Wild edible plant Diversity of Kotagiri hills- a part of Nilgiri Biosphere Reserve, Southern India, Journal of Research in Biology, 2011; 2: 80-87.
- Yeshodharan K, Sujana KA, Wild edible plants traditionally used by the tribes in the Parambikulam Wildlife sanctuary, Kerala, India, Natural Product Radiance, 2007; 6 (1): 74-80.
- Kar A, Common wild vegetables of Aka tribe of Arunachal Pradesh, Indian journal of Traditional Knowledge, 2004; 3(3): 305-313.
- Binu S, Wild edible plants used by the tribals in pathanamthitta district, Kerala, Indian Journal of Traditional knowledge, 2010; 9 (2): 309-312.
- Rajasab AH, Isaq M, Documentation of Knowledge on edible wild plants of North Karnataka, Indian Journal of Traditional Knowledge, 2004; 3(4): 419-429.
- Kayang H, tribal knowledge on wild edible plants of Meghalaya, Northeast, India, Indian Journal of Traditional Knowledge, 2007; 6 (1): 177-181.
- Ramachandran VS, Wild edible plants of Anamalais, Coimbatore district, Western Ghats, Tamil Nadu, Indian Journal of Traditional Knowledge, 2007; 6 (1): 173-176.
- Patil MV, Some more Wild edible plants of Nashik district (Maharashtra), Ancient science of life, 2000; XIX (3 & 4): 102-104.

19. Angami A, Gujrel PR, Rethy P, Singh B, Kalitha SK, Status and potential of wild edible plants of Arunachal Pradesh, Indian Journal of Traditional Knowledge, 2006; 5(4): 541-550.
20. Sinha R, Lakra V, Edible weeds of tribals of Jharkhand, Orissa and West Bengal, Indian Journal of Traditional Knowledge, 2007; 6 (1): 217- 222.
21. Kala CP, Prioritization of cultivated and wild edible by local people in the Uttaranchal hills of Indian Himalaya, Indian Journal of Traditional Knowledge, 2007; 6 (1): 239- 244.
22. Jain AK, Tiwari P, Nutritional value of some traditional edible plants used by tribal communities during emergency with reference to Central India, Indian Journal of Traditional Knowledge, 2012; 11(1): 51-57.
23. Arinathan V, Mohan VR, De Britto AJ, Mrugan C, wild edibles used by Palliyars of the western Ghats, Tamil Nadu, Indian Journal of Traditional Knowledge, 2007, 6 (1): 163-168.
24. Valvi SR, Deshmukh SR, Rathod VS, Ethnobotanical survey of wild edible fruits in Kolhapur district, International Journal of Applied Biology and Pharmaceutical Technology, 2011; 2 (1): 194-197.
25. Pal M M, Janifer R X, Phani Kumar G, Gupta S & Shashi B S, Phytofoods of Nubra Valley, Ladakh- The Cold desert, Indian Journal of Traditional Knowledge, 2010, 9 (2): 303- 308.
26. Ajesh TP, Abdulla Naseef SA & Kumuthakalavalli R, Ethnobotanical documentation of wild edible fruits used by Muthuvan Tribes of Idukki, Kerala- India, International Journal of Pharma and Biosciences, 2012; 3 (3): 479-487.
27. Reddy KN, Pattanaik C, Reddy CS, Raju VS, Traditional Knowledge on wild food plants in Andhra Pradesh, Indian Journal of Traditional Knowledge, 2007; 6 (1): 223-229.
28. Rethy P, Singh B, Kagyung R, Gajurel PR, Ethnobotanical studies of Dehang- Debang Biosphere reserve of Arunachal Pradesh with special reference to Memba Tribe, Indian Journal of Traditional Knowledge, 2010; 9 (1): 61-67.
29. Jain A, Singh H B & Bhattacharya P R, The Ethnobotany and Nutritional values of wild rice [*Zizania latifolia* (Griseb.) Turcz. ex Stapf] (Poaceae) in Manipur, Indian Journal of Traditional Knowledge, 2012; 11(1): 66-69.
30. Singh B, Sinha BK, Phukan SJ, Borthakur SK, Singh VN, Wild edible plants used by Garo tribes of Nokrek Biosphere Reserve in Meghalaya, India, Indian Journal of Traditional Knowledge, 2012; 11 (1): 166-171.
31. Chandra S, Saklani S, Dimari M, Evaluation of Garhwal Himalaya Wild edible tuber *Dioscorea deltoidea*, International Research Journal of Pharmacy, 2012; 3 (3): 152-156.
32. Chanra S, Saklani S, Singh J, Phytochemical screening of Garhwal Himalaya Wild edible tuber *Colocasia esculenta*, Int Res J Pharm 2012; 3 (3): 181-186.
33. Naik VN. Flora of Marathwada. Vol. 1. (Ranunculaceae to Convolvulaceae). Amrut Prakashan Aurangabad, 1998.
34. Naik VN. Flora of Marathwada. Vol. 2. (Solanaceae to Poaceae). Amrut Prakashan Aurangabad, 1998.
35. Yadav SR, Sardesai MM. Flora of Kolhapur District. Shivaji University Kolhapur Maharashtra, India, 2002.
36. Yadav SR.,Kahalkar VI., Bhuskute SM. A new species of *Amorphophalus* Bl. Ex. Decne. (Araceae) From Bhandara district, Maharashtra State, India. Aroideana. 2009; 32: 132-135.
37. Kahalkar V. I., Floristic study on Gondia district of Maharashtra State. Ph.D. thesis, Nagpur University Nagpur, 2009.

Cite this article as:

Patale Chandrakumar K, Nasare Praveenkumar N, Narkhede Sushama D. Ethnobotanical studies on wild edible plants of Gond, Halba and Kavar tribes of Salekasa taluka, Gondia district, Maharashtra state, India. Int. Res. J. Pharm. 2015; 6(8):512-518
<http://dx.doi.org/10.7897/2230-8407.068103>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.