Research Article

www.ijrap.net

EPIDEMIOLOGICAL STUDY OF KLAIBYA IN HYPERTENSIVE PATIENTS

Pandey Anurag^{1*}, Tiwari Mamta², Godatwar Pawankumar¹, Sevatkar B.K.¹

¹Department of Rog and Vikriti Vijnana, National Institute of Ayurveda, Jaipur, Rajasthan, India

²Department of Swasthavritta and Yoga, National Institute of Ayurveda, Jaipur, Rajasthan, India

Received on: 20/05/12 Revised on: 23/06/12 Accepted on: 19/08/12

*Corresponding author

Dr. Anurag Pandey, PG Scholar, Department of Rog and Vikriti Vijnana, National Institute of Ayurveda, Jaipur, Rajasthan, India.

E-mail: dr.anuragpandey@yahoo.in

DOI: 10.7897/2277-4343.03645

Published by Moksha Publishing House. Website www.mokshaph.com

All rights reserved.

ABSTRACT

Klaibya (Male Sexual dysfunction) is the most burning problem among the sexual diseases because in this condition the person is unable to perform the coitus due to complete or partial lack of erection and/or rigidity and if coitus attempted it ends into failure and dissatisfaction. The incidence of Klaibya is increasing day by day with increase in the incidence of diabetes, Hypertension, peripheral vascular disorders, peripheral neuropathy, anxiety, stress, depression and their medications.

Male sexual dysfunction is more prevalent in Hypertensive's than Normotensive individuals, and several mechanisms have been implicated in the pathogenesis of sexual dysfunction in hypertensive patients. Sexual dysfunction represents a major quality-of-life related health problem, and available data indicate that Hypertension is a major risk factor for Klaibya (Male sexual dysfunction) in men.

In the present study, the survey was carried out on 1000 subjects of general society to access the prevalence of Klaibya with and without hypertension.

Key Words- Klaibya, Male Sexual dysfunction, IIEF-15, JNC-7

INTRODUCTION

Male Sexual Dysfunction or "Klaibya¹" can be defined as a "man's inability to attain or maintain an erection of sufficient strength to perform the act of intercourse²" "Patient should seek medical advice if this occurs consistently over 6 month period and in more than 50% of attempts"³. The basic aetiology behind this include psychological, neurological, vascular insufficiency, drug side effects as antihypertensive drugs, anti depressants drugs, diseases such as Diabetes Mellitus and others.

Sexual Dysfunction defined by the World Health Organization as "the various ways in which an individual is unable to participate in a sexual relationship as he or she would wish⁴". As it is obvious, sexual dysfunction affects both men and women. Sexual dysfunction has been considered of psychogenic origin since ancient times; however, it is currently believed to be a disease of organic (mainly vascular) etiology in the majority of cases. Since atherosclerosis of the arteries supplying genital tissues greatly affects sexual function, it seems rational to assume that conditions predisposing to atherosclerosis (hypertension, diabetes, obesity, hyperlipidemias) might impair sexual function. Blood pressure is a major contributor in the atherosclerotic process and vascular and per vascular genital tissues exhibit profound damage with increasing blood pressure⁵.

Aims and objectives

This research work aimed to conduct an Epidemiological (Nidanatmaka) study of Klaibya and establish the etiopathological significance of Hypertension in pathogenesis of Klaibya (Male sexual dysfunctions).

For this work, survey based case record form were prepared on the basis of compilations incorporating Dosha, Dushyadi factors as well as IIEF-15 and JNC-VII

criteria. Survey was conducted at World Ayurvedic Congress Camp Bangalore, National Institute Ayurveda Jaipur and Local camps outside Jaipur, where data was collected to find out the prevalence in general society.

MATERIALS AND METHODS

Inclusion criteria

- Adult male patients (above 18 years of age) suffering from Klaibya (Male Sexual Dysfunctions) with or without Hypertension (IIEF-15⁶ and JNC-VII⁷).
- Adult male patients (above 18 years of age) suffering from Hypertension with or without Klaibya (Male Sexual Dysfunctions) (IIEF-15 and JNC-VII).

IIEF-15 Variables⁸

DOMAIN 1-

Erectile Function in study population (n=1000).
Que.no.1,2,3,4,5 and 15.

Que No 1- Frequency of erection during sexual activity
Que No 2-Frequency of erection hard enough for penetration

Que No 3- Frequency of able to penetrate (enter)

Que No 4- Frequency of maintain erection after penetrated (entered)

Que No 5- Frequency of maintain erection to completion of intercourse

Que No 15- Frequency of confidence that can get and keep erection

DOMAIN 2-

Intercourse Satisfaction in study population (n=1000).
Que no.6,7, and 8.

Que No 6- how many times have attempted sexual intercourse

Que No 7- Frequency of satisfaction

Que No 8- Frequency of enjoyed sexual intercourse

DOMAIN 3-

Orgasmic Function in study population (n=1000).

Que no.9 and 10.

Que No 9- Frequency of ejaculation

Que No 10- Frequency of feeling of orgasm or climax (with or without ejaculation)

DOMAIN 4 –

Sexual Desire in study population (n=1000).

Que no.11 and 12.

Que No 11- Frequency of sexual desire

Que No 12- Frequency of level of sexual desire

DOMAIN 5-

Overall Satisfaction in study population (n=1000).

Que no. 13 and 14.

Que No 13- Frequency of satisfaction with overall sex life

Que No 14- Frequency of satisfaction with sexual relationship with partner.

JNC –VII guidelines⁹

Category	SBP mm/Hg		DBP mm/Hg
Normal	<120	and	<80
Prehypertension	120–139	or	80–89
Hypertension, Stage 1	140–159	or	90–99
Hypertension, Stage 2	>160	or	>100

Exclusion Criteria

- Female patients.
- Malignant Hypertension.¹⁰
- Patients having mechanical cause of Klaibya (Male Sexual Dysfunctions) like Antisperm antibodies condition, Obstruction, Surgical process.

Institutional Ethics Committee Permission

As per provisions, before initiation of the project the protocol submitted to the ethics committee for approval. The information and consent form also submitted to the ethics committee on the date 07-04-2010. The (CTRI) Clinical Trials Registry- India number for this trial is REF/2012/06/3686.

RESULTS

Salient observations of Epidemiological study

Table 1: Showing the prevalence of Klaibya and Hypertension in study population (n=1000)

Sr. No.	Condition	Number of subjects	Percentage
Klaibya	Klaibya without Hypertension	392	39.2%
	Klaibya with Hypertension	248	24.8%
	Total Klaibya	640	64%
Hypertension	Hypertension without Klaibya	206	20.6%
	Hypertension with Klaibya	248	24.8%
	Total Hypertension	454	45.5%
People without Klaibya or Hypertension		154	15.4%

Table 2: Age wise distribution of 1000 Subjects

Age Groups (Years)	People those who are affected by Klaibya or Hypertension				People those who are not affected by Klaibya or Hypertension	Total
	Klaibya	Klaibya with Hypertension	Hypertension	Total		
18 up to 30	79	39	19	137	78	215
31 up to 43	172	92	48	312	32	344
44 up to 57	93	64	66	223	28	251
58 up to 70	48	53	73	174	16	190
Total				846	154	1000

- Klaibya (n= 392)
- Klaibya + hypertension (n = 247)
- Hypertension (n = 206)
- People without Klaibya or Hypertension (n = 154)

Figure 1: Depicting the prevalence of Klaibya and Hypertension in a group of volunteers (n=1000)

Figure 2: Depicting the Marital Status wise distribution (n= 1000)

Figure 3: Depicting the Education wise distribution (n=1000)

Figure 4: Depicting the Prevalence of Hypertension in study population (n=1000)

Figure 5: Depicting the Prevalence of Klaibya in study population (n=1000)

Figure 6: Depicting the Difference between self Assessment and IIEF-15 Assessment wise distribution

ANALYSIS OF IIEF-15 VARIABLES

Showing the prevalence of DOMAIN 1- Erectile Function in study population (n=1000). Que.no. 1, 2,3,4,5 and 15

Figure 7: Depicting the Prevalence of DOMAIN 1 - Erectile Functioning in study population (n=1000)

Showing the prevalence of DOMAIN 2- Intercourse satisfaction in study population (n=1000). Que no. 6, 7, and 8

Figure 8: Depicting the Prevalence of DOMAIN 2 Intercourse satisfaction in study population (n=1000)

Showing the prevalence of DOMAIN 3- Orgasmic function in study population (n=1000). Que no. 9 and 10

Figure 9: Depicting the Prevalence of DOMAIN 3 Orgasmic functioning in study population (n=1000)

Showing the prevalence of DOMAIN 4- Sexual desire in study population (n=1000). Que no. 11 and 12

Figure 10: Depicting the Prevalence of DOMAIN 4 Sexual Desire in study population (n=1000)

Showing the prevalence of DOMAIN 5- Overall Satisfaction in study population (n=1000). Que no. 13 and 14

Figure 11: Depicting the Prevalence of DOMAIN 5 Overall Satisfaction in study population (n=1000)

DISCUSSION

In this Survey, study of Klaibya with or without Hypertension revealed that the prevalence of Klaibya and Hypertension in 1000 population, 64% suffered from Klaibya and 45 % suffered from Hypertension. Among 248 people two groups suffered from both Klaibya and hypertension, while only a meager 15.4 % (n=154) people were not affected by either Klaibya or hypertension.

In the prevalence of Domain-1 maximum number of people i.e. 344 suffer from Moderate dysfunction followed by 324 mild dysfunction, 173 were of severe dysfunction, 129 were of no dysfunction and 30 were of mild to moderate dysfunction.

In the prevalence of Domain -2 maximum number of people 264 suffered from no intercourse dissatisfaction followed by 229 moderate intercourse dissatisfaction, 224 were of mild intercourse dissatisfaction, 213 were of Mild to moderate intercourse dissatisfaction and 70 were of severe intercourse dissatisfaction.

In the prevalence of Domain -3 maximum number of people 268 suffered from mild dysfunction followed by 250 moderate dysfunction, 195 were of mild to, moderate

dysfunction, 146 were of severe dysfunction and 141 were of no dysfunction.

In the prevalence of Domain -4 maximum number of people 259 suffer from Mild to moderate dysfunction followed by 252 mild dysfunction, 251 were of moderate dysfunction, 179 were of severe dysfunction and 59 were of no dysfunction.

In the prevalence of Domain - 5 maximum number of people 376 suffer from Mild to moderate dysfunction followed by 229 moderate dysfunctional, 228 were of mild dysfunction, 100 were of severe dysfunction and 67 were of no dysfunction.

SUMMARY

In this study a survey was conducted on 1000 volunteer as having symptoms related to Hypertension and Klaibya and result showed that significant number of people suffered from Klaibya (n = 640) while 454 people suffered with hypertension. In 248 people two groups suffered from both Klaibya and hypertension. Only a meager 15.4 % (n=154) people were not affected by either Klaibya or hypertension.

CONCLUSION

On the basis of survey based study we can conclude that Hypertension and Klaibya have strong relation with each other's pathogenesis. In our study we found that person suffering from Klaibya were found to have Hypertension and the persons having Hypertension also suffered with Klaibya. Most of the person having Hypertension was not aware with Klaibya; it may be due to social stigma burden attached with this disease (Klaibya). They were found under the disease when using the IIEF-15 Variables. So on the basis of our study we can conclude that each patient of Hypertension should be examined for Klaibya.

ACKNOWLEDGEMENT

I feel great pleasure in expressing my sincere gratitude and deep sense of indebtedness to my respected Guide Dr. Pawan Kumar Godatwar, Associate Professor of P.G. department of Roga and Vikriti Vijnana N.I.A, without whom working on this subject would not have been materialized. His ever inspiring, helpful attitude and parental affection enabled me to pursue this study with continuous zeal.

ABBREVIATIONS

MSD- Male Sexual Dysfunctions

SD- Sexual Dysfunctions

IIEF- International Index of Erectile Function

JNC- Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure

SBP: Systolic Blood Pressure

DBP: Diastolic Blood Pressure

REFERENCES

1. Charak Samhita With Ayurveda Dipika Commentary Of Chakrapani Datta, Ed. Yadavji Trikarma Ji Acharya, Chaukhambha Surbharti Prakashan, Varanasi. Edition 7th, 2002, Chikitsa Sthana Chapter 30 Verse 155-157, pp 641 Smith's, General Urology, 12th edition, pp 666
2. Roger S Kirby, An Atlas of Erectile Dysfunction, Second edition, published in the Taylor and Francis e-Library, 2005
3. Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States; prevalence and predictors. JAMA 1999;281:537-44 <http://dx.doi.org/10.1001/jama.281.6.537>
4. Grimm RH, Grandits GA, Prineas RJ, et al. Long-term effects on sexual function of five antihypertensive drugs and nutritional hygienic treatment in hypertensive men and women: Treatment of Mild Hypertension Study (TOMHS). Hypertension 1997;29:8-14. <http://dx.doi.org/10.1161/01.HYP.29.1.8> PMID:9039073
5. Rosen RC, Riley A, Wagner G, Osterloh IH, Mishra A. The international index of erectile function (IIEF): a multidimensional scale for assessment of erectile dysfunction: Journal of Urology. 1997 Jun; 49(6): 822-30. [http://dx.doi.org/10.1016/S0090-4295\(97\)00238-0](http://dx.doi.org/10.1016/S0090-4295(97)00238-0)
6. Chobanian AV, National High Blood Pressure Education Program Coordinating Committee. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 report. JAMA 2003 May 21; 289:2560-72. <http://dx.doi.org/10.1001/jama.289.19.2560> PMID:12748199
7. Rosen RC, Riley A, Wagner G, Osterloh IH, Kirkpatrick J, Mishra A., IIEF. International Index of Erectile Function Questionnaire.
8. Chobanian AV, National High Blood Pressure Education Program Coordinating Committee. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. The JNC 7 report. JAMA 2003 May 21; 289:2560-74 <http://dx.doi.org/10.1001/jama.289.19.2560> PMID:12748199
9. Thomas L. Managing hypertensive emergencies in the ED. Can Fam Physician 2011;57 (10):1137-97 PMID:21998228 PMCid:3192077

Cite this article as:

Pandey Anurag, Tiwari Mamta, Godatwar Pawankumar, Sevatkar BK. Epidemiological study of Klaibya in hypertensive persons. nt J Res Ayur & Pharm. 2012; 3(5):888-892

Source of support: Nil, Conflict of interest: None Declared