Review Article

www.ijrap.net

CATHARANTHUS ROSEUS (TAPAK DARA): "A CONTROVERSIAL" MEDICINAL PLANT IN INDONESIA

Em Sutrisna*

Senior lecturer, Department of Pharmacology of Faculty of Medicine of Universitas Muhammadiyah Surakarta, Indonesia

Received on: 04/06/15 Revised on: 23/07/15 Accepted on: 02/08/15

***Corresponding author**

Dr. EM Sutrisna, MD, Senior lecturer, Department of Pharmacology of Faculty of Medicine of Universitas Muhammadiyah Surakarta, Indonesia.
Email: Em.Sutrisna@ums.yahoo.com and em_sutrisna@yahoo.com

DOI: 10.7897/2277-4343.065117

ABSTRACT

Herbal medicine industry in Indonesia was shaken when Indonesian National Agency of Drugs and Food control prohibits the development of herbal medicines made from *Catharanthus roseus*. The reason given is that *Catharanthus roseus* contains alkaloids causing bone marrow depression. The purpose of this article to explore the effects of both preclinical and clinical pharmacological of *Catharanthus roseus*. The result of the research shows *Catharanthus roseus* have many pharmacological effects include: antidiabetic, hypolipidemic, antihypertension, anticancer, antiangiogenesis, antispermatogenic, antifungal, antibacteria, antioxidant, wound healing and even hepatoprotective activity.

Keywords: *Catharanthus roseus*, Herbal medicine, Pharmacological, Alkaloid, Bone suppression

INTRODUCTION

Indonesian National Agency of Drugs and Food Control released regulation no. 10 of 2014 that prohibit the development (manufacture & distribute) of herbal medicines / supplements made from *Catharanthus roseus*. This raises questions and allegations as to be *Catharanthus roseus* is a dangerous plant.¹ The regulation stated that the alkaloids in *Catharanthus roseus* may cause bone marrow depression. Interestingly, many studies suggest a variety of pharmacological effects of *Catharanthus roseus*. Most studies are still in the preclinical stage study, but some others already in early stages of clinical trials. Vincristine and Vinblastin of *Catharanthus roseus* even been marketed. The prohibition should not be necessary if the development of herbal medicines made from *Catharanthus roseus* can eliminate the presence of alkaloids. This review tries to investigate pharmacological effects of *Catharanthus roseus*.

Description

Catharanthus roseus is tropical/subtropics plant that is spread throughout the world.² This plant belongs to the family of Apocynaceae. This plant is herbaceous/ sub herbaceous with a high CMM 40-80. The flower color are purple and pink white.³ It has different names in several countries, among others: *Tapak dara* (Indonesia), *Kemuning china* (Malaysia), *Periwinkle* (Madagascar), *Nayantara* (Bangladesh), *Ainskati*, *Nityakalyani*, *Rattan jot*, *Sadaphul*, *Ushamanjairi* (India), *Kantotan* (Filipina), *Nichinich-so* (Jepang), *Chatilla* (Guatemala), *Phaeng phoi farang* (Thailand), etc.

Traditional Uses

Leaves of *Catharanthus roseus* is commonly used as antidiabetic in Northern Europe (Swanston-Flatt) and

India.⁴ In Brazil, Dominica, England, Cook Islands, Jamaica, Mozambique, Pakistan, Taiwan, Thailand and West Indies, decoction of *Catharanthus roseus* leaves are also used as antidiabetic.⁵ In India this plant was used to treat depression, muscle pain, bleeding gums, mouth, ulcers and sore throats.⁶ This plant was also used to treat cystitis, gastritis, hypertension, enteritis, diarrhea.⁷ In South Africa, people use *Catharanthus roseus* as urogenital infection,⁸ diabetes mellitus (Zulu people),⁹ menorrhagia¹⁰ and rheumatism.¹¹ In Southern and Eastern Africa, this plant was used to treat unspecified venereal diseases.¹² The people of Mutirikwi, Zimbabwe, used *Catharanthus roseus* to treat stomach ache.¹³ The decoction of leaves of *Catharanthus roseus* were used to treat diabetes mellitus (In the Kancheepuram, a District of Tamil Nadu, India).¹⁴

Preclinical studies of pharmacological properties

Antidiabetic

The ethanolic extract of *Catharanthus roseus* leaves & flower reduced blood glucose level in diabetic rats.^{15,16} The combination of dichloromethane : methanol extract with ratio 1:1 dosage 500mg/kg bw orally for 7 days and 15 days reduced blood glucose level with hypoglycemic activity 48.6 and 57.6% in streptozotocin induced diabetic rats.¹⁷ *Catharanthus roseus* leaf powder suspension in 2 ml distilled water (100 mg/kg body weight/day/60 days) orally can reduced plasma glucose and increase plasma insulin in diabetic rat induced by streptozotocin.¹⁸ The aqueous extract of *Catharanthus roseus* leaves dose 1g/kgbw can reduced blood glucose level significantly as compared to control (28.20±0.34 vs 32.99±0.89 m mol/L).¹⁹ The aqueous extract of *Catharanthus roseus* leaves, roots, flower and stem at doses 250mg/kgbw intra peritoneal for 14 days can reduced blood glucose level and the reduction percentage is 52,90%.²⁰ The combination of

dichloromethane : methanol (1:1) extract doses 500mg/kgbw orally 20 days can decrease blood glucose level in male wistar albino rats compared to control diabetic (110.08 ± 8.6 vs 241.62 ± 2.3 mg/dL).²¹ The juice of *Catharanthus roseus* leaf dose 0.5; 0.75 and 1.0ml/kgbw orally can decrease blood glucose level after 6, 8, 10, 12, 16, 18, 20 & 24 h drug administration with reduction percentage of 16.7% (6 h, $p < 0.05$), 28.6% (18 h, $p < 0.05$), and 31.9% (20 h, $p < 0.01$).²² The combination of dichloromethane: methanol (1:1) extract of whole plant of *Catharanthus roseus* (500mg/kg body weight) daily orally for 14 days can reduced blood glucose level on diabetic female albino rats wistar strain induced by alloxan 80 mg/kgbw.²³ The aqueous extract of *Catharanthus roseus* can decreased blood glucose on diabetic rats significantly.²⁴ Research by Islam *et al.*, found that the ethyl acetate fraction of ethanolic extract of *Catharanthus roseus* leaves dose of 150 mg/kgbw intra peritoneal can reduce fasting blood glucose on 24 h with 40.6 percent of reduction on diabetic rats induced by streptozotocin (45 mg/kg).²⁵ The chemical compound of this fraction that hypoglycemic effect are alkaloid like as catharanthin, leurosine, lochnerine, tetrahydroalstonin, vindoline and vindolinine ($p < 0.01$).²⁶

Hypolipidemic

The *C. Roseus* leaf powder suspension in 2 ml distilled water (100 mg/kgbw/day) for 60 days orally can decrease total cholesterol, triglyceride, LDL, VLDL, atherogenic index but it increases HDL.¹⁸ The aqueous extract of *Catharanthus roseus* of dose of 250 mg, 350 mg, and 450 mg/kg body weight for 30 days can reduce lipid profile (TC, TG, LDL & VLDL) significantly in diabetic rats.²⁴ The fraction of petroleum ether, ethyl acetate and chloroform of ethanolic extract of *Catharanthus roseus* leaves at dose of 150 mg/kgbw intraperitoneally can decrease total cholesterol, triglyceride after 24 h.²⁵

Antispermaticogenic

The aqueous extract of *Catharanthus roseus* has antispermaticogenic on the male Swiss albino mice. The mice were treated by this extract had a sperm count and percentage of motile sperm less than control. The dead sperm of treated mice was more than control.²⁷

Antihypertensive

Research by Ara *et al.*, 2009 found that the leaves extract of *Catharanthus roseus* at dose of 30 mg/155±15 g body weight by intra peritoneal injection for one week reduced blood pressure on hypertensive rats induced by adrenalin. This effect was equivalent with atenolol at dose of 0.1mg/155±15 g body weight.²⁸

Antifungal

The ethanolic extract of *Catharanthus roseus* leaf has antifungal activity toward *Aspergillus niger*, *Aspergillus flavus*, *Aspergillus fumigates*, *Candida albicans* and *Penicillium chrysogenum* by disc diffusion method. The width of zone of inhibition 4.2 mm; 3mm; 1.8mm; 1.6mm and 3.3mm respectively. Secondary metabolites such as alkaloids in *Catharanthus roseus* were suspected causing antifungal effect.²⁹

Anticancer and cytotoxic

The chloroform extracts of crude of *Catharanthus roseus* has cytotoxic activity to human colorectal carcinoma cell line (HCT 116). Vindolin and catharanthine are two isolated compound from *Catharanthus roseus* which has cytotoxic activity to HCT 116. The cytotoxic activity of catharanthine was greater than vindolin. Catharanthine has cytotoxic activity to HCT 116 with IC_{50} 60µg/mL.³⁰ Vincalukoblastine derivative from alkaloid vinea has antineoplastic activity in mice, was transplanted by leukemia L1210, P1534 and Ehrlich ascites tumor cells,³¹ while vincristine has antineoplastic effect on patients with Hodgkin's disease. 39% patients with Hodgkin's disease treated by vincristine have temporary remission more than one month.³² The mechanism involved were binding of this agent to tubulin. This binding could block the mitosis cell that causes metaphase capture.^{33,34}

Antibacterial

The 10, 25, 50, 75 and 100 mg of the dried whole plant extracts with dichloromethane: methanol (1:1) have antibacterial activity to *K. pneumoniae*, *P. aeruginosa*, *E. coli*, *B. cereus*, *B. subtilis*, and *S. aureus*.²³ Research by Balaabirami *et al.*, 2012 found that the ethanolic extract of *Catharanthus roseus* leaves has antibacterial effect to *Escherichia coli*, *Klebsiella oxytoca*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *Salmonella paratyphi* and *Staphylococcus aureus*.²⁹

Hepatoprotective

The ethanolic extract of *Catharanthus roseus* leaves at dose of 500 mg/kgbw reduced blood SGOT and SGPT level in rats. This effect may be due to by antioxidant property of phenolic compound from this extract.³⁵

Antioxidant

The methanolic extract of *Catharanthus roseus* val. Alba leaves at doses of 200 & 400 mg/kgbw have antioxidant effect. This extract reduced plasma superoxide dismutase, catalase and glutathione levels significantly of diabetic rats as compared to controls.³⁶

Wound healing activity

The ethanolic extract of *Catharanthus roseus* flower 100mg/kgbw/day orally for 10 days showed wound healing activity in wound model rats SD strain. This extract increased the wound breaking strength in the incision wound and decreased wound area and epithelization period in excision wound compared with controls ($P < 0.001$).³⁷ Research by Dewi *et al.*, 2013 found that the methanol leaf extract of *Catharanthus roseus* 15% topically increases wound healing/epithelization in wistar rats.³⁸

Antiangiogenesis

Several plants of traditional Chinese medicinal herb have anti-angiogenesis effect on chick embryo chorioallantoic membrane model. The aqueous extract of *Catharanthus roseus* leaves at dose of 1g herb/mL has anti-angiogenesis activity with percentage of inhibition is ($33.65 \pm 1.62\%$).³⁹

Clinical studies of pharmacological properties**Anticancer**

Vincristine sulfate is indicated to treat lymphoblastic leukemia acute and lymphoma. This agent was approved by US FDA in 1963 with brand name Oncovin®. Vinblastine is indicated to treat lymphomas inclusive of Hodgkin's disease, bladder and breast cancers. In 1965, Vinblastine sulfate was approved by FDA. This agent was marketed with brand name (Velban®).^{34,40}

Vindesine and Vinorelbine, a semisynthetic derivate of vinblastine have potency to treat hematologic malignancy. Vindesine (Eldisine®) is used in France, the UK, and other countries to treat drug-resistant acute of lymphoid leukemia,⁴¹ while vinorelbine shows potency to treat Hodgkin's disease.^{42,43} Vinorelbine tartrate (Navelbine®) is used to treat non-small cell lung carcinoma or advanced breast cancer.⁴⁰

Antidiabetic

In a clinical study involved 60 people suffered from diabetic mellitus, the administration of the aqueous extract of *Catharanthus roseus* at dose of 75mg/day for 30 days can reduce blood glucose level 2 h post prandial, glycated Hb, Total Cholesterol, Total Triglyceride, LDL-C, VLDL-C, HDL-C significantly as compared to control (placebo).⁴⁴

Circulatory disorder

Ajmalicine, alkaloid from *Catharanthus roseus*, is used for treatment circulatory disease.⁴⁵ The monomeric alkaloids ajmalicine and serpentine are used in the treatment of circulatory diseases.⁴⁶⁻⁵³

Chemical compounds

Catharanthus roseus contains about 130 different alkaloids.⁵⁴ The several alkaloid among others: reserpine, ajmalicine, vinceine, raubasin,⁵⁴ catharantine and vindoline.³⁰ Anticancer drugs are vincristine & vinblastine,³⁰ cartharathine, lochnerine, vindoline, vindolinine, tetrahydroalstronine, reserpinne, serpentine.²⁶ Vincristine is indicated to treat Hodgkin's Lymphoma, while vinblastine revealed to treat leukemia on child,⁵⁵ yohimbin,⁵⁶ and flavonoid hirsutidin.⁵⁷

The Phytoconstituents of *Catharanthus roseus* are carbohydrate, alkaloid, glycoside, flavonoids, tannin, saponins, protein, amino acid, fats and oils.²³ Fraction of petroleum ether of ethanolic of *C. roseus* leaves contains tannin, alkaloid, flavonoid, while fraction of chloroform & ethyl acetate contains alkaloid and flavonoid.²⁵

THE PROFILE OF HERBAL MEDICINE INDUSTRY IN INDONESIA

Indonesia is a tropical country and known as having second largest biodiversity after Brazil. It is estimated that there are around 25000-30000 plants found in Indonesia. This quantity is about 80% of the total populations plants in the world and 90% of the population of plants in Asia.^{58,59} It is estimated that 7000 species of plants are utilized by Indonesian people as a traditional medicine and about 283 species of medicinal herbs are used by industry and registered to Indonesia National Agency of Drugs and Food Control.⁵⁹

The development of herbal medicine industry in Indonesia continues to increase. In 2002 recorded only 29 small traditional medicine industry. This number increased to 172 in 2006. The complete list can be seen in Table 1

Table 1: Herbal Medicine industry profile in Indonesia.⁶⁰

Year	Small industry of Traditional medicine	Industry of Traditional medicine	Pharmacy industry	Total
2002	29	10	16	55
2003	164	58	82	304
2004	217	54	85	356
2005	197	47	87	331
2006	172	40	79	291

Research by the Agency for Health Research and Development of Ministry of Health in 2007 on the use of herbal medicine by the people of Indonesia showed that 35.7% of people use herbs and more than 85% of them admitted the beneficial herbs for health.⁶¹ In 2010 research was repeated and the result were 59.12% and 95.6% people admitted the beneficial herbs for health.⁶²

Only 32 herbal medicine industries in Indonesia have been following good manufacturing practices for herbal medicine.⁶³ The majority of herbal medicine in Indonesia are not concerned with the alkaloids content of *Catharanthus roseus* that might cause bone marrow depression. This possibly underlies the prohibition of this herb production by Indonesia National agency of Food and Drugs Control.

CONCLUSION

Catharanthus roseus has many pharmacological effects. The prohibition of the development of herbal medicines made from *Catharanthus roseus* due to its alkaloids contents could be avoided if the industry can eliminate these alkaloids. Within certain limits this rule can be understood as the practical level. Most Indonesian herbal medicine industries are small and medium industries which are sometimes less concern with the risk of alkaloids resulted from the *Catharanthus roseus*.

REFERENCES

- Indonesia National agency of Drugs and Food control. Peraturan kepala badan pengawas obat dan makanan Republik Indonesia. Jakarta. 2014. No. 10.pp.1-4.
- Karthikeyan B., Jaleel CA, Changxing Z, Joe MM, Srimannarayan J, Deiveekasundaram. M. The effect of AM fungi and phosphorous level on the biomass yield and ajmalicine production in *Catharanthus roseus*. Eur.Asia J BioSci. 2008;2: 26-33
- Padua, de LS,Bunyapraphatsara N and Lemmens RHMJ. Plant sources of shouthes Asia. Prosea foundation, Bogor. Indonesia 1999;12:411-412
- Tiong SH, Looi CY, Hazni H, Arya A, Paydar M, Wong WF, Cheah SC, Mustafa MR and Awang K. Antidiabetic and Antioxidant Properties of Alkaloids from *Catharanthus roseus* (L.) G. Don. Molecules 2013; 18: 9770-9784 <http://dx.doi.org/10.3390/molecules18089770>
- Don G. *Catharanthus roseus*, Medicinal plants of the world. Human Press, Totowa, NJ, 1999.pp.109-118
- Gajalakshmi S, Vijayalakshmi S and Rajeswari DV. Pharmacological activities of *Catharanthus roseus*:a perspective review. Int J Pharm Bio Sci. 2013;4(2): 431 – 439
- Dessisa D. Preliminary economic evaluation of medicinal plants in Ethiopia: trade,volume and price In: Medhin Z,Abebe D, editors.

- Proceedings of the National Workshop on biodiversity Conservation and Sustainable use of Medicinal Plants in Ethiopia. Addis Ababa, Ethiopia: 28 th April-May1 st , 2001; 176-188
8. Fernandes I, Van Rensburg CEJ, Hoosen AA, Steenkamp V. In vitro activity of medicinal plants of the Venda region, South Africa, against *Trichomonas vaginalis*. S. Afr. J. Epidemiol. Infect. 2008; 23:26-28
9. Dauskardt R. The changing geography of traditional medicine, urban herbalism on the Witwatersrand, South Africa. Geogr. J. 1990; 22:257-283 <http://dx.doi.org/10.1007/bf00711339>
10. Hutchings A, Scott AH, Lewis G, Cunningham AB. Zulu medicinal plants-Aninventory. University of Natal Press, Pietermaritzburg. 1996.
11. Marles RJ & Farnsworth NR. Antidiabetic plants and their active constituents. Phytomedicine.1995; 2:137-189 [http://dx.doi.org/10.1016/S0944-7113\(11\)80059-0](http://dx.doi.org/10.1016/S0944-7113(11)80059-0)
12. Watt JM, Breyer-Brandwijk MG. The Medicinal and Poisonous Plants of Southern and Eastern Africa. Second Edition, Livingstone, London.1962 pp.25-100
13. Chigora P, Masocha R, Mutenheri F. The role of indigenous medicinal knowledge (IMK) in the treatment of ailments in rural Zimbabwe: The case of Mutirikwi communal lands. J. Sustain. Dev. Afr. 2007; 9:1520-5509
14. Muthu C, Ayyanar M, Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. J. Ethnobiol. Ethnomed 2006; 2:1-10 <http://dx.doi.org/10.1186/1746-4269-2-43>
15. Ghosh RK & Gupta I. Effect of *Vinca Rosea* and *Ficus racemosus* on hyperglycemia in rats . Indian J. Anim Health. 1980;19:145-149
16. Chattopadhyay RR, Sarkar RK, Ganguly S, Banerjee RN, Basu TK. Hypoglycemic and anti hyperglycemic effect of leaves of *Vinca rosea* Linn. Ind J physiol pharmacol. 1991; 35:145-151
17. Singh SN, Vats P, Suri S, Shyam R, Kumria MM, Ranganathan S, Sridharan K. Effect of an antidiabetic extract of *Catharanthus roseus* on enzymic activities in streptozotocin induced diabetic rats. J. Ethnopharmacol. 2001;76(3):269-77. [http://dx.doi.org/10.1016/S0378-8741\(01\)00254-9](http://dx.doi.org/10.1016/S0378-8741(01)00254-9)
18. Rasineni K, Bellamkonda R, Singareddy SR, and Desireddy S. Antihyperglycemic activity of *Catharanthus roseus* leaf powder in streptozotocin-induced diabetic rats. Pharmacognosy Res. 2010; 2(3): 195-201. <http://dx.doi.org/10.4103/0974-8490.65523>
19. Mostofa, M. Choudhury E, Hossain MA, Islam HM, Islam MS and Sumon MH. Antidiabetic effects of *Catharanthus roseus*, *Azadirachta indica*, *Allium sativum* and glimepride in experimentally diabetic induced rat. Bangl. J. Vet. Med.2007;5(1&2): 99-102
20. Ávila EV, Cano-Velasco JL, Alarcón-Aguilar FJ, Ortiz MCJ, Almanza-Pérez JC, and Román-Ramos R. Hypoglycemic Activity of Aqueous Extracts from *Catharanthus roseus*. Evidence-Based Complementary and Alternative Medicine. Volume 2012: 1-7
21. Jayanthi M, Sowbala N, Rajalakshmi G, Kanagavalli U, and Sivakumar V. Study of anti hyperglycemic effect of *Catharanthus roseus* in alloxan induced diabetic rats. International Journal of Pharmacy and Pharmaceutical Sciences. 2010;2(4): 114-116
22. Nammi S, Boini MK, Lodagala SD and Behara RBS. The juice of fresh leaves of *Catharanthus roseus* Linn. reduces blood glucose in normal and alloxan diabetic rabbits. BMC Complementary and Alternative Medicine. 2003; 3: 1-4 <http://dx.doi.org/10.1186/1472-6882-3-1>
23. Ibrahim M, Mehjabeen SS, and Narsu ML. Pharmacological evaluation of *Catharanthus roseus*. International Journal of Pharmaceutical Applications. 2011; 2(3):165-173
24. Natarajan A , Ahmed KSZ , Sundaresan S, Sivaraj A , Devi K, & Kumar BS. Effect of aqueous flower extract of *Catharanthus roseus* on alloxan induced diabetes in male Albino rats . International Journal of Pharmaceutical Sciences and Drug Research 2012; 4(2): 150-153
25. Islam MA, Afia Akhtar M, Khan MRI, Hossain MS, Alam, MK, Wahed MII, Rahman BM, Anisuzzaman ASM, Shaheen SM, and Ahmed M . Antidiabetic and Hypolipidemic Effects of Different Fractions of *Catharanthus Roseus* (Linn.) on Normal and Streptozotocin-induced Diabetic Rats . J. Sci. Res. 2009;1 (2):334-344
26. Gordon, S. H., Marvin, G. and Marry, R. A. Alkaloids of *Vinca rosea*; A Preliminary report on hypoglycemic activity. Lloydia.1964; 27:361-363
27. Murugavel TA, Ruknudin ST, Akbarsha MA. Antifertility effect of *Vinca rosea* (Linn.) leaf extract on male albino mice – a sperm parametric study. Curr Sci.1989;58(19): 1102- 1103
28. Ara N, Rashid M and Amran S .Comparison of hypotensive and hypolipidemic effects of *catharanthus roseus* leaves extract with atenolol on adrenaline induced hypertensive rats. Pak. j. Pharm. Sci. 2009;22(3): 267-271
29. Balaabirami S and Patharajan S. In vitro antimicrobial and antifungal activity of *Catharanthus roseus* leaves extract against important pathogenic organisms. Int J Pharm Pharm Sci, 2012; 4 (3): 487-490
30. Siddiqui MJ, Ismail Z, Aisha AFA & Abdul majid AMS, Cytotoxic activity of *C roseus* (Apocynaceae) crude extract and pure compound against human colorectal carcinoma cell line. Int J.o Pharmacology.2010; 6(1): 43-47
31. Cutts JH, Beer CT, Noble RL. Biological properties of vincalkebblastine, an alkaloid in *Vinca rosea* Linn. with reference to its antitumor action. Cancer Res. 1960; 20:1023-1031.
32. Bohannon RA, Miller DG and Diamonds HD. Vincristine in the Treatment of Lymphomas and Leukemias. Cancer Research.1963; 23: 613-620
33. Cutts JH. The effect of vincalkebblastine on dividing cells *in vivo*. Cancer Res.1961;21:168-172
34. Chabner BA, Amrein PC, Druker BJ, In: Goodman & Gilman's The Pharmacological Basis of Therapeutics. 11th Edition. Brunton LL, Lazo JS, Parker KL, editors. New York: McGraw-Hill; 2006. pp. 1257-1262.
35. Ahmed MF & Rao A S. Comparative Hepatoprotective Activities of Selected Indian Medicinal Plants. Global Journal of Medical research Pharma, Drug Discovery, Toxicology and Medicine.2013;13(2):15-21
36. Barkat A & Mujeeb M. The comparative study of *Catharanthus roseus* extract and extract loaded chitosan nanoparticles in alloxan induced diabetics rats. Int.j.Bio.Res.2013;4(12):670-678
37. Nayak B, Pereira LMP. *Catharanthus roseus* flower extract has wound-healing activity in Sprague Dawley rats. BMC Complementary and Alternative Medicine. 2006; 6(41): 1-6
38. Dewi IALP, Damriyasa IM, Dada IKA. Bioaktivitas Ekstrak Daun Tapak Dara (*Catharanthus Roseus*) Terhadap Periode Epitelisasi Dalam Proses Penyembuhan Luka Pada Tikus Wistar. Indonesia Medicus Veterinus. 2013; 2(1) : 58 – 75
39. Wang S, Zheng Z, Weng Y, Yu Y, Zhang D, Fan W, Dai R, Hu Z. Angiogenesis and anti-angiogenesis activity of Chinese medicinal herbal extracts. Life Sciences 2004;74: 2467 – 2478 <http://dx.doi.org/10.1016/j.lfs.2003.03.005>
40. Guéritte F, Fahy J. In: Anticancer Agents from Natural Products. Cragg GM, Kingston DGI, Newman DJ, editors. Boca Raton, FL: CRC/Taylor & Francis; 2005. pp. 123-135.
41. Dancey J, Steward WP. The role of vindesine in oncology--recommendations after 10 years' experience. Anticancer Drugs. 1995; 6:625-636. <http://dx.doi.org/10.1097/00001813-199510000-00001>
42. Devizzi L, Santoro A, Bonfante V, Viviani S, Balzarini L, Valagussa P *et al*. Vinorelbine: an active drug for the management of patients with heavily pretreated Hodgkin's disease. Ann Oncol.1994;5:817-820.
43. Diehl V, Re D, Harris NL, P.M. M. In: Cancer: Principles and Practice of Oncology.8th Ed. DeVita VT, Lawrence TS, Rosenberg SA, editors. Philadelphia: Williams & Wilkins; 2008. pp. 2167-2220.
44. Bisla G, Choudhary S, Singh E and Chaudhary V . Hyperglycemia and hyperlipidemia mitigating impact of *Catharanthus roseus* (*Sadabahar*) leaves aqueous extract on type2 diabetes mellitus subjects. Asian Journal of Plant Science and Research. 2013, 3(4):170-174
45. Misra N, Luthra R, Kumar S. Enzymology of indole alkaloid biosynthesis in *Catharanthus roseus*. Indian J Biochem Biophys.1996;33(4):261-73
46. Jaleel CA, Gopi R, Alagu Lakshmanan GM, Panneerselvam R. Triadimefon induced changes in the antioxidant metabolism and ajmalicine production in *Catharanthus roseus* (L.) G. Don. Plant Science. 2006;171: 271-276. <http://dx.doi.org/10.1016/j.plantsci.2006.03.018>
47. Jaleel CA, Gopi R, Manivannan P, Panneerselvam R. Antioxidative potentials as a protective mechanism in *Catharanthus roseus* (L.) G. Don. plants under salinity stress. Turkish Journal of Botany.2007a; 3: 245-251.
48. Jaleel CA, Gopi R, Manivannan P, Panneerselvam R. Responses of antioxidant defense system of *Catharanthus roseus* (L.) G. Don. to paclobutrazol treatment under salinity. Acta Physiologiae Plantarum.2007b; 29: 205-209. <http://dx.doi.org/10.1007/s11738-007-0025-6>

49. Jaleel CA, Gopi R, Sankar B, Manivannan P, Kishorekumar A, Sridharan R, Panneerselvam R. Studies on germination, seedling vigour, lipid peroxidation and proline metabolism in *Catharanthus roseus* seedlings under salt stress. South African Journal of Botany.2007c; 73: 190-195. <http://dx.doi.org/10.1016/j.sajb.2006.11.001>
50. Jaleel CA, Manivannan P, Kishorekumar A, Sankar B, Panneerselvam R. Calcium chloride effects on salinity induced oxidative stress, proline metabolism and indole alkaloid accumulation in *Catharanthus roseus*. Comptes Rendus Biologies.2007d; 330(9): 674-683. <http://dx.doi.org/10.1016/j.crv.2007.07.002>
51. Jaleel CA, Manivannan P, Sankar B, Kishorekumar A, Gopi R, Somasundaram R, Panneerselvam R. Induction of drought stress tolerance by ketoconazole in *Catharanthus roseus* is mediated by enhanced antioxidant potentials and secondary metabolite accumulation. Colloids and surfaces B: Biointerfaces. 2007e; 60(2): 201-206. <http://dx.doi.org/10.1016/j.colsurfb.2007.06.010>
52. Jaleel CA, Manivannan P, Sankar B, Kishorekumar A, Sankari S, Panneerselvam R. Paclitaxel enhances photosynthesis and ajmalicine production in *Catharanthus roseus*. Process Biochemistry. 2007f; 42: 1566-1570. <http://dx.doi.org/10.1016/j.procbio.2007.08.006>
53. Jaleel CA, Panneerselvam R. Variations in the antioxidative and indole alkaloid status in different parts of two varieties of *Catharanthus roseus*, an important folk herb. Chinese Journal of Pharmacology and Toxicology. 2007g; 21(6): 487-494.
54. Sandeep P, Jagjit K, Raman K & Kuldeep K. *Catharanthus roseus*: a medicinal plant with potent anti tumor properties. Int.J. Res.Ayurveda Pharm.2014;5(6):653-654
55. Johnson IS, Armstrong JG, Gorman M and Burnett JP. The vinca alkaloids : A new class of oncolytic agent. Cancer Res. 1963;23:1390-1427
56. Millan, MJ, Newman TA, Audinot V, Cussac D, Lejeune F, Nicolas JP, Cogé F, Galizzi JP and Boutin JA. Agonist and antagonist actions of Yohimbine as compared to fluparoxan at alpha (2)- adrenergic receptors (AR). Synapse. 2002;35: 79-95 [http://dx.doi.org/10.1002/\(SICI\)1098-2396\(200002\)35:2<79::AID-SYN1>3.0.CO;2-X](http://dx.doi.org/10.1002/(SICI)1098-2396(200002)35:2<79::AID-SYN1>3.0.CO;2-X)
57. Piovan, A. and R. Fillipini. Anthocyanins in *Catharanthus roseus* in vivo and in vitro: A review. Phytochem. Rev. 2007; 6: 235-242. <http://dx.doi.org/10.1007/s11101-006-9052-y>
58. Erdelen WR, Adimihardja K, Moesdarsono H, Sidik. Biodiversity, traditional medicine and the sustainable use of indigenous medicinal plants in Indonesia. Indigenous knowledge and development monitor. 1999; 7(3):3-6.
59. Pramono E. The commercial use of traditional knowledge and medicinal plants in Indonesia. Submitted for multi-stakeholder dialogue on trade, intellectual property and biological resources in Asia, 2002.
60. Direktorat Penilaian Obat Tradisional, Suplemen Makanan dan Kosmetik–Badan POM, Jakarta.2007.
61. Indonesian Health Ministry. Riset Kesehatan Dasar.Jakarta. 2007.
62. Indonesian Health Ministry. Riset Kesehatan Dasar.Jakarta. 2010
63. Oktavia, R & Maharesi, R.2012. Sistem Penunjang Keputusan Penjadwal Proses Produksi Industri Jamu Berdasarkan Good Manufacturing Practices (GMP). http://repository.gunadarma.ac.id/1139/1/SISTEM%20PENUNJANG%20KEPUTUSAN_UG.pdf. accessed April 2015

Cite this article as:

Em Sutrisna. *Catharanthus roseus* (Tapak dara):”a controversial” medicinal plant in Indonesia. Int. J. Res. Ayurveda Pharm. 2015;6(5):629-633 <http://dx.doi.org/10.7897/2277-4343.065117>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IJRAP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJRAP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IJRAP editor or editorial board members.