Research Article

IDENTIFICATION OF MAIZE LINE OF MANIPUR CONTAINING *crtRB1* 3'TE FAVOURABLE ALLELE USING MOLECULAR MARKER

YAIKHOM VIVEKANANDA*¹, WAREPAM JESMI DEVI², JADUMANI AHONGSHANGBAM¹, SINGH N.B.¹ AND LAISHRAM J.M.¹

¹Department of Genetics & Plant Breeding, College of Agriculture, Central Agricultural University, Imphal, 795004, India

²Department of Biotechnology, H.K. School of Life Sciences, Assam University, Silchar, 788011, India

*Corresponding Author: Email - boneybi@gmail.com

Received: May 19, 2018; Revised: May 26, 2018; Accepted: May 27, 2018; Published: May 30, 2018

Abstract- Vitamin A is one of the essential nutrients for normal functioning and growth in our body. Biofortification of vitamin A in the crop (Maize) is an economical way to overcome vitamin A deficiency. The *crtRB1* is an important gene in the carotenoid pathway which increases β -carotene accumulation in maize. Selecting the lines having favourable allele is a cost-effective method alternative to high performance liquid chromatography (HPLC). In the present study, we screened hundred maize lines of Manipur derived from CAU-87 using a *crtRB1* 3'TE gene-specific marker and found that three lines viz., CAU-87-9, CAU-87-12, CAU-87-28 contained the favourable allele which can be utilized for maize improvement programme.

Key words- Maize, β -carotene, *crtRB1* gene, Manipur

Citation: Yaikhom Vivekananda, et al., (2018) Identification of Maize Line of Manipur Containing *crtRB1* 3'TE Favourable Allele Using Molecular Marker. International Journal of Genetics, ISSN: 0975- 2862 & E-ISSN: 0975-9158, Volume 10, Issue 5, pp.-434-437.

Copyright: Copyright©2018 Yaikhom Vivekananda, et al., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Academic Editor / Reviewer: Ch. Dhananjay Singh

Introduction

Vitamin A or Retinol is one of the important essential nutrient required in our body for normal growth and functioning [1]. World Health Organization (WHO) reported that due Vitamin A Deficiency (VAD) alone, globally around 0.25 to 0.5 million children become blind every year and half of them died within 12 months of losing sight. Since our body cannot synthesize Vitamin A of its own, it needs to be provided as provitamin A, carotenoids, including α -carotene, β -carotene and β -cryptoxanthin [2] which further converted into vitamin A through metabolism. One of the sustainable and cost-efficient method to overcome vitamin A deficiency is through biofortification of β -carotene into food crops. Maize is one of the important food crops and together with rice and wheat, it accounts at least 30% of the food calories to more than 4.5 billion people [3]. It is also one of the selected crops for vitamin A biofortification, which is one of the world's most widely grown cereal crop and is ranked third among cereal [4]. Biofortification of vitamin A in Maize is an economical way to overcome VAD, globally. [2,5]. The four predominant carotenoids found in maize are β -carotene, β -cryptoxanthin, zeaxanthin and lutein. However, β -carotene has the most provitamin A activity among the carotenoid found [6,7]. Recent research in maize has demonstrated three important genes in the accumulation of provitamin A in grain viz., Phytoene synthase1 (Y1 or *Psy1*) which catalyses the formation of phytoene and responsible for the yellow colour of grains [8]; *lcyE* (lycopene epsilon cyclase) which converts lycopene to α -lycopene with association of other gene and natural occurring mutant allele of this gene with less functionality divert the more lycopene towards β -carotene branch enhancing more flux towards provitamin A accumulation; *crtRB1* gene (β -carotene hydroxylase 1) an important gene associated with β -carotene accumulation which converts β -carotene (having 100% provitamin A activity) to β -cryptoxanthin (having 50% pro-vitamin A activity) and zeaxanthin (having 0% pro-vitamin A activity). The naturally occurring variation of this gene can increase more accumulation of β -carotene by preventing further conversion into other

compounds [1,2,7,9]. Quantifying of provitamin A contain using high performance liquid chromatography (HPLC) is costly and time consuming; marker-assisted selection (MAS) using DNA markers can help in identifying the genotypes with high accumulation of provitamin A. [10,11]. *Psy1* can increase β - carotene accumulation by increasing the quantity of substrate in the carotenoid biosynthesis pathway. The favourable allele of *Psy1* is nearly fixed in the tropical maize and favourable alleles of *lcyE* and *crtRB1*, however, are uncommon in tropical germplasm and may be targeted for high provitamin A [12] maize breeding in the tropical region. The *crtRB1* had a huge and significant impact on enhancing β -carotene and total provitamin A content, regardless of genetic constitution for *lcyE*- 5'TE and Marker Assisted Selection (MAS) for favourable 'allele 1' of *crtRB1* can lead to speedy doubling or more of total provitamin A concentration while MAS for favourable 'allele 4' of *lcyE* normally results in 20-30 % increase in total provitamin A concentration. Three polymorphisms of *crtRB1* have been identified through mapping approach viz., 5'TE (in the 5'- Untranslated Region), InDel4 (in the coding region) and 3'TE (spanning the sixth exon and 3'- Untranslated Region) and PCR based co-dominant markers were identified for all the three polymorphisms helping in improvement of provitamin A content in maize through Marker Assisted Selection [1,2]. The 3'TE polymorphism of *crtRB1* gene (that spans the 6th exon and The 3'TE polymerism of *crtRB1* gene has three alleles that have a varying degree of β -carotene accumulation in maize kernel viz. allele 1 without TE insertion (543 bp), allele 2 having 325 bp insertion (296 bp +875 bp) and allele 3 having 1250 bp insertion (296 bp+1221 bp+1880 bp). The allele 1 (favourable allele) enhanced the β -carotene concentration by reducing transcript expression of the *crtRB1* gene, with an average increase of 6.50 μ g/g β -carotene in maize endosperm comparing with unfavourable allele (allele 2 and allele 3) [1, 2]. The North-Eastern Himalayan region of India including the state of Manipur is rich in maize diversity which is supported by the Asiatic origin of maize [13].

Fig-1 Allelic variation of 3' TE of *crRB1* gene in maize lines of Manipur. Lane M- 200bp DNA ladder; Lane 1 to Lane 12- line CAU-87-1 to CAU-87-12. Lane 9, 12- lines with favourable allele *i.e.*, CAU-87-9, CAU-87-12.

Table-1 Allelic polymorphism of 3'TE of *crRB1* gene in Maize lines of Manipur

SN	Lines(Progeny from CAU87)	Allelic size(bp)	Sl. No.	Lines(Progeny from CAU87)	Allelic size (bp)
1.	CAU-87-1	1221+296	51.	CAU-87-51	1221+296
2.	CAU-87-2	1221+543+296	52.	CAU-87-52	543+296
3.	CAU-87-3	1221+296	53.	CAU-87-53	1221+296
4.	CAU-87-4	1221+296	54.	CAU-87-54	1221+543+296
5.	CAU-87-5	1221+296	55.	CAU-87-55	1221+296
6.	CAU-87-6	1221+296	56.	CAU-87-56	1221+296
7.	CAU-87-7	1221+543+296	57.	CAU-87-57	296
8.	CAU-87-8	1221+543+296	58.	CAU-87-58	1221+296
9.	CAU-87-9	543	59.	CAU-87-59	1221+296
10.	CAU-87-10	1221+543+296	60.	CAU-87-60	296
11.	CAU-87-11	1221+296	61.	CAU-87-61	296
12.	CAU-87-12	543	62.	CAU-87-62	543+296
13.	CAU-87-13	543+296	63.	CAU-87-63	296
14.	CAU-87-14	1221+543+296	64.	CAU-87-64	543+296
15.	CAU-87-15	1221+296	65.	CAU-87-65	1221+543+296
16.	CAU-87-16	1221+296	66.	CAU-87-66	1221+543+296
17.	CAU-87-17	1221+296	67.	CAU-87-67	296
18.	CAU-87-18	1221+296	68.	CAU-87-68	543+296
19.	CAU-87-19	1221+543+296	69.	CAU-87-69	543+296
20.	CAU-87-20	1221+543+296	70.	CAU-87-70	1221+543+296
21.	CAU-87-21	1221+296	71.	CAU-87-71	543+296
22.	CAU-87-22	1221+296	72.	CAU-87-72	1221+296
23.	CAU-87-23	1221+296	73.	CAU-87-73	543+296
24.	CAU-87-24	1221+543+296	74.	CAU-87-74	1221+543+296
25.	CAU-87-25	1221+296	75.	CAU-87-75	543+296
26.	CAU-87-26	1221+543+296	76.	CAU-87-76	1221+296
27.	CAU-87-27	1221+543+296	77.	CAU-87-77	1221+296
28.	CAU-87-28	543	78.	CAU-87-78	1221+296
29.	CAU-87-29	1221+296	79.	CAU-87-79	296
30.	CAU-87-30	1221+296	80.	CAU-87-80	296
31.	CAU-87-31	1221+296	81.	CAU-87-81	296
32.	CAU-87-32	543+296	82.	CAU-87-82	296
33.	CAU-87-33	1221+296	83.	CAU-87-83	296
34.	CAU-87-34	1221+296	84.	CAU-87-84	296
35.	CAU-87-35	296	85.	CAU-87-85	296
36.	CAU-87-36	1221+296	86.	CAU-87-86	1221+296
37.	CAU-87-37	296	87.	CAU-87-87	296
38.	CAU-87-38	1221+296	88.	CAU-87-88	296
39.	CAU-87-39	543+296	89.	CAU-87-89	1221+296
40.	CAU-87-40	543+296	90.	CAU-87-90	1221+543+296
41.	CAU-87-41	543+296	91.	CAU-87-91	1221+296
42.	CAU-87-42	296	92.	CAU-87-92	1221+296
43.	CAU-87-43	296	93.	CAU-87-93	1221+296
44.	CAU-87-44	296	94.	CAU-87-94	1221+296
45.	CAU-87-45	296	95.	CAU-87-95	1221+296+543
46.	CAU-87-46	296	96.	CAU-87-96	1221+296
47.	CAU-87-47	543+296	97.	CAU-87-97	1221+296
48.	CAU-87-48	296	98.	CAU-87-98	543+296
49.	CAU-87-49	296	99.	CAU-87-99	296
50.	CAU-87-50	296	100.	CAU-87-100	543+296

Fig-2 Allelic variation of 3' TE of *crtRB1* gene in maize lines of Manipur. Lane M- 200bp DNA ladder; lane 1 to lane 12- line CAU-87-38 to CAU-87-49; lane 13, 14- line CAU-87-28 and CAU-87-32. Lane 13-line with favourable allele *i.e.*, CAU-87-28.

Recent genotyping of maize cultivars of Manipur [14] have found that CAUCH 87 or CAU 87 landrace of Manipur contained favourable allele (543 bp) of *crtRB1* 3' TE gene in heterozygous condition with unfavourable allele (296 bp). Since the presence of 296 bp (unfavourable allele with the insertion of 325 bp in 3' TE region) can lower the β -carotene accumulation. The present study aims at selecting the lines or progeny derived from CAU-87 landrace having favourable allele 1 (543 bp).

Materials and Method

Plant Materials

A hundred seeds (kernels) randomly selected from seed lot of self-population of CAU 87 were planted in the laboratory and later transplanted in the field. Each plant was assigned with specific code for the study. The lines or progeny (from the single kernel) derived from CAU 87 were screened for the presence of favourable allele.

DNA extraction

DNA was extracted from the leaves of 15 days old seedlings using SDS procedure with slight modification [15].

PCR analysis

Screening for favourable allele 1 (543bp) was done using *crtRB1* 3' TE gene-specific primers, *crtRB1*-3' TE-F: ACACCACATGGACAAGTTCG, *crtRB1*-3' TE-R1: ACACTCTGGCCCATG AACAC and *crtRB1*-3' TE-R2: ACAGCAATACAGGGGACCAG [2,7,10]. The primers were synthesized and purified by IDT. PCR was performed with a total volume of 25 μ L, containing 1 μ L crude DNA, 2.5 μ L of PCR buffer (supplied with *Taq* polymerase), 0.2 mM each of dNTPs, 0.3 μ M of each primer and 2 U of *Taq* DNA Polymerase (GeNei, Bengaluru, India). PCR amplification (2720 Thermal Cycler, Applied Biosystems, California, USA) was carried out using the standard cycle conditions [7]. The PCR products were separated by electrophoresis on 2% agarose gel. The gels were stained with ethidium bromide and detected under Gel Doc XR+ with Image Lab Software (Bio-Rad, California, USA).

Results and Discussion

Out of the hundred lines screened for the presence of a favourable allele of *crtRB1* 3' TE gene, three lines *i.e.*, CAU-87-9, CAU-87-12 and CAU-87-28 have found to contain the favourable allele 1 (543 bp) in homozygous condition. Thirty-two lines have 543 bp in heterozygous conditions (16 lines having 543bp +296bp allele and 16 lines having 1221 bp+543 bp+296 bp allele) which can be further self for recovering progeny having favourable (543bp) in homozygous condition. Remaining progeny or lines have an unfavourable allele of 3' TE gene *i.e.*, either 296 bp or 1221+296 bp which is given in table 1. Presence of favourable allele of

crtRB1 3' TE alone can double the β -carotene content [2] and the line having that allele can be utilized for marker-assisted backcross for improvement vitamin A content in maize breeding programme.

Conclusion

The present genotyping of maize lines of Manipur for the presence of a favourable allele (543 bp) of *crtRB1* 3' TE found that three lines *i.e.*, CAU-87-9, CAU-87-12 and CAU-87-28 derived from CAUCH 87 have the favourable allele and can be utilized for marker-assisted backcross programme for improvement of vitamin A content in maize.

Application of research: Improvement of maize for vitamin A content.

Research Category: Genetics and Plant Breeding

Abbreviations:

%: Percentage
 μ g: Microgram
 μ L: Microlitre
 μ M: Micromolar
 3' TE: 3' Transposable element
 3' UTR: Untranslated region
 bp: Base pair
crtRB1: β - carotene hydroxylase 1
 DNA: Deoxyribonucleic Acid
 dNTP: Deoxyribonucleotide Triphosphate
 g: Gram
 HPLC: High Performance Liquid Chromatography
 IDT: Integrated DNA Technologies
lcyE: Lycopene epsilon cyclase
 MAS: Marker Assisted Selection
 mM: Millimolar
 Psy 1 or Y1: Phytone synthase 1
 SDS: Sodium dodecyl sulphate
 U: unit
 VAD: Vitamin A Deficiency
 WHO: World Health Organisation

Acknowledgement / Funding: Author thankful to Central Agricultural University, Imphal, 795004, India

*Research Guide or Chairperson of research: JM Laishram
 University: Central Agricultural University, Imphal, 795004, India

Research project name or number: PhD Thesis

Author Contributions: All author equally contributed

Author statement: All authors read, reviewed, agree and approved the final manuscript

Conflict of Interest: No Conflict of Interest

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

References

- [1] Vignesh M., Hossain F., Nepolean T., Choudhary M., Saha S., Bhat J.S., Prasanna M.B., Gupta H.S. (2012) *Indian J. Genet.*, 72,189–194.
- [2] Yan J.B., Kandianis C.B., Harjes C.E., Bai L., Kim E., Yang X.H., Skinner D., Fu Z.Y., Mitchell S., Li Q., Fernandez M.G.S., Zaharieva M., Babu R., Fu Y., Palacios N., Li J.S., DellaPenna D., Brutnell T., Buckler E.S., Warburton M.L., Rocheford T. (2010) *Nat Genet.*, 2(4),322-7.
- [3] Shiferaw B., Prasanna B. M., Hellin J., Banziger M. (2011) *Food Security*,3, 307-327.
- [4] Ayisi K.K., Poswall M.A.L. (1997) *Europ. J. Agron.*, 16,151–159.
- [5] Tanumihardjo S.A., Bouis H., Hotz C., Meenakshi J.V., McClafferty B. (2008) *Compr. Rev. Food Sci. Food Safety*,7, 329–334.
- [6] Harjes E.C., Rocheford R.T., Ling B., Brutnell P.T., Kandianis B.C., Sowinski G.S., Stapleton E.A., Vallabhaneni R., Williams M., Wurtzel T.E., Yan J., Buckler S.E. (2008) *Sci.*, 319, 330–333.
- [7] Selvi T.D., Senthil N., Yuvaraj N., John Joel A., Mahalingam A., Nagarajan P., Vellaikumar S., Srimathi P., Raveendran M. and Nepolean T. (2014) *Food Biotechnology*, 28, 41-49.
- [8] Li Q., Farre G., Naqvi S., Breitenbach J., Sanahuja G., Bai C., Sandmann G., Capell T., Christou P., Zhu C. (2010) *Transgenic Res.*, 19,1053–1068.
- [9] Babu R., Rojas N.P., Gao S., Yan J., Pixley K. (2012) *Theor. Appl. Gene.*, 126, 389–399.
- [10] Muthusamy V., Hossain F., Thirunavukkarasu N., Choudhary M., Saha, S. (2014) *PLoS ONE*, 9(12), 1-22.
- [11] Zunjare R.U., Hossain F., Muthusamy V. (2017) *Plant Breed.*, 136, 872–880.
- [12] Fu Z., Chai Y., Zhou Y., Yang X., Warburton M.L., Xu S., Cai Y., Zhang D., Li J., Yan J. (2013) *TheorApplGenet*, 126, 923–935.
- [13] Anjula P., Semwal D.P., Ahlawat S.P., Sharma S.K. (2015) *Maize (Zea mays): Collection Status, Diversity Mapping and Gap Analysis, National Bureau of Plant Genetic Resources, New Delhi, India*, 34.
- [14] Chuwanghijam, Dhananjoychingangbam, Thangjamsurchandra, Laishram J.M. (2017) *International Journal of Agricultural Science and Research*, 7(1),115-120.
- [15] Zheng K., Subudhi P.K., Domingo J., Magpantay G. and Huang N. (1995) *Rice Genetics Newsletter*, 12, 255–258.